



# Family A GPCR heteromers in animal models

Javier González-Maeso \*

Departments of Psychiatry and Neurology, Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA

\*Correspondence: [javier.maeso@mssm.edu](mailto:javier.maeso@mssm.edu)

**Edited by:**

Dominique Massotte, Institut des Neurosciences Cellulaires et Intégratives, France

**Reviewed by:**

Ralf Jockers, University of Paris, France

**Keywords:** G protein-coupled receptor (GPCR), dimerization, heteromer, heterocomplex, serotonin 5-HT<sub>2A</sub> receptor, metabotropic glutamate 2 receptor, molecular pharmacology

## INTRODUCTION

G protein-coupled receptors (GPCRs) were assumed to exist and function in the plasma membrane as monomeric proteins that became activated by binding of one agonist ligand to one receptor molecule (Bourne et al., 1990). However, although previous findings based on rather indirect measures such as radioligand binding had suggested a direct interaction of two receptors with each other (Limbird et al., 1975; Ferre et al., 1991), it was the application of a protein-protein interaction assay by bioluminescence resonance energy transfer (BRET) that revealed the phenomenon of molecular proximity between beta2-adrenergic receptors in living cells (Angers et al., 2000). Since then, this topic has been a major subject of research and numerous *in silico* and *in vitro* studies have suggested expression of family A GPCRs as homodimers and higher-order homomers in heterologous expression systems. However, the demonstration that reconstitution of a single beta2-adrenergic receptor molecule into lipoprotein particles leads to efficient activation of G proteins raised concerns about the functional significance of family A GPCR homomers (Whorton et al., 2007), and this is currently a controversial topic (for an extensive review on GPCR homodimers/homomers, see Milligan, 2013; see also Bouvier and Hebert, 2014; Lambert and Javitch, 2014).

Another fundamental yet relatively independent question is that related to expression of different GPCR subtypes as heteromers. It is well accepted that the family C GABA<sub>B</sub> receptor needs two protomers (GABA<sub>B</sub>-R1 and GABA<sub>B</sub>-R2) to reach the plasma membrane as a functional dimeric receptor (Jones et al.,

1998; Kaupmann et al., 1998; White et al., 1998). On the other hand, although multiple lines of evidence indicate that family A GPCR heteromers may exist, particularly in tissue cultures (González-Maeso, 2011; Ferre et al., 2014), only relatively recent studies started to test this formulation in whole animal models.

## FAMILY A GPCR HETEROMERS IN WHOLE ANIMAL MODELS

One of the main limitations of the classical techniques used to define GPCR heteromeric formation is the translation of findings obtained in cellulo into physiological or behavioral assays in whole animal models. In this context, co-immunoprecipitation is an approach commonly used to examine protein-protein interaction in native tissue (Milligan and Bouvier, 2005). GPCR antibodies are usually neither specific nor sensitive and therefore validation assays in knockout mice are often required (Fribourg et al., 2011; Moreno et al., 2012). Considering this, it is also clear that demonstration of co-immunoprecipitation in native tissues does not imply the existence of a heteromeric assembly, as they may form part of same protein complex through for example PDZ domain-binding motifs at the end of the C-terminal tails of both receptor types together with adaptor proteins (Magalhaes et al., 2010). Remarkably, there are only a few studies that have investigated GPCR heteromeric formation in living animals, and due to the lack of biophysical methods applicable to study protein-protein interactions in preclinical models, their experimental approaches were mostly focused on signaling and behavioral outcomes rather than on the

existence of molecular proximity between different GPCR subtypes.

Although it does not measure molecular proximity, an attractive approach to define whether heteromeric formation is involved in behavioral phenotypes is the use of peptides that disrupt receptor complex formation. These peptides tested *in vivo* are usually selected according to findings previously obtained in heterologous expression systems. As an example, it was demonstrated that the G<sub>s</sub>-coupled dopamine D<sub>1</sub> receptor and the G<sub>i/o</sub>-coupled dopamine D<sub>2</sub> form a receptor complex that induces Ca<sup>2+</sup> release via a G<sub>q/11</sub>-dependent pathway (Lee et al., 2004), and that the region of Met-257—Glu-271 (intracellular loop 3; D2<sub>IL3-29-2</sub>) but not Asn-243—Ile-256 (intracellular loop 3; D2<sub>IL3-29-1</sub>) of the dopamine D<sub>2</sub> receptor can pull-down the dopamine D<sub>1</sub> receptor. Based on a Tat-tagged peptide approach, it was shown that intracerebroventricular administration of the peptide D2<sub>IL3-29-2</sub>, which disrupts heteromeric formation between dopamine D<sub>1</sub> and D<sub>2</sub> receptors *in vitro*, induces antidepressant-like effects in rats (Pei et al., 2010). More recent findings using serial deletions and point mutations further demonstrate that dopamine D<sub>1</sub> receptor carboxyl tail residues Glu-404 and Glu-405 are critical in mediating the interaction with the D<sub>2</sub> receptor, and that administration of a disrupting peptide Tat-D1 modulates depression-like behavior in rats such as forced swim test (Hasbi et al., 2014). A similar approach was used to block the association as a GPCR heteromer between the mu-opioid receptor isoform MOR1D and the gastrin-releasing peptide receptor (GRPR) in the spinal cord (Liu et al., 2011). The authors demonstrated

that the C-terminus of MOR1D is critical for MOR1D-GRPR heteromeric formation. Using a Tat-fusion peptide, they also found that a motif consisting of seven amino acids of the MOR1D C-terminus (RNEEPSS) attenuates morphine-induced scratching, but not morphine-induced analgesia.

The question of whether GPCR heteromers exist *ex vivo* has been addressed using time-resolved Förster resonance energy transfer FRET (TR-FRET) in plasma membrane preparations of mouse brain. It was found that the dopamine D<sub>2</sub> receptor and the ghrelin receptor (GHSR1a) co-localize in mouse striatum, hippocampus and hypothalamus (Kern et al., 2012). When membrane preparations from hypothalamus were incubated with red-ghrelin (acceptor fluorophore) and an anti-D<sub>2</sub> receptor antibody together with a europium cryptate-labeled secondary antibody (donor fluorophore), a significantly TR-FRET signal was observed. Although TR-FRET signal is eliminated in hypothalamic membrane preparations of GHSR1a knockout mice, which supports specificity, these findings were observed *ex vivo* in plasma membrane preparations and further investigation will be necessary to confirm the existence of GHSR1a-D<sub>2</sub> heteromeric formation in hypothalamus *in vivo*.

Another indirect approach to test whether GPCR heteromeric formation affects behavioral phenotypes is the use of chimeric constructs that according to biophysical assays in tissue culture do not form heteromeric complexes. Examples include the 5-HT<sub>2A</sub>-mGlu2 heteromeric receptor complex (Gonzalez-Maeso et al., 2008) and the MT<sub>1</sub>-MT<sub>2</sub> melatonin heteromeric receptor complex (Baba et al., 2013). Serotonin 5-HT<sub>2A</sub> and metabotropic glutamate 2 (mGlu2) receptors have been shown to form a GPCR heteromeric complex in HEK293 cells. Using chimeric constructs, it was demonstrated that three residues located at the intracellular end of TM4 of mGlu2 are necessary to form a complex with the 5-HT<sub>2A</sub> receptor (Ala-677<sup>4,40</sup>, Ala-681<sup>4,44</sup>, Ala-685<sup>4,48</sup>) (Fribourg et al., 2011; Moreno et al., 2012). Head-twitch is a rodent behavior model induced by hallucinogenic 5-HT<sub>2A</sub> agonist such as lysergic

acid diethylamide (LSD) and DOI (Hanks and Gonzalez-Maeso, 2013). This behavior requires expression of 5-HT<sub>2A</sub> receptor in cortical pyramidal neurons (Gonzalez-Maeso et al., 2007) and is absent in mGlu2 knockout mice (Moreno et al., 2011), which supports that mGlu2 is necessary for 5-HT<sub>2A</sub>-dependent behavioral events. Using a virally-mediated (HSV) over-expression approach, it was demonstrated that the head-twitch response induced by the hallucinogenic 5-HT<sub>2A</sub> receptor agonist DOI was rescued in mGlu2 knockout mice over-expressing wild-type mGlu2 in frontal cortex, and that this did not occur in mGlu2 knockout mice over-expressing mGlu2deltaTM4N—a mGlu2/mGlu3 chimeric construct that according to previous findings *in vitro* and in cellulo does not form the 5-HT<sub>2A</sub>-mGlu2 receptor heteromer (Moreno et al., 2012). A similar approach was used to investigate function of the MT<sub>1</sub>-MT<sub>2</sub> melatonin heteromeric receptor complex *in vivo* in mouse (Baba et al., 2013). The electroretinogram (ERG), consisting mainly of an a-wave and a b-wave, is commonly used to assess retinal function. Using transgenic mice that express MT<sub>2</sub>-P95L (mutant that does not form the MT<sub>1</sub>-MT<sub>2</sub> heteromeric receptor complex in HEK293 cells), it was shown that control mice responded to melatonin injection with an increase in the amplitude of the a-wave and b-wave, whereas MT<sub>2</sub>-P95L did not.

Although these events have been proposed to represent a demonstration of GPCR heteromeric expression, thereby suggesting a new target for drug design, their conclusions in animal models were based largely on indirect approaches that measured phenotypes affected by manipulations such as chimeric constructs or Tat-tagged peptides that impact heteromeric organization *in vitro*. Consequently, it remains unclear as to whether different GPCR subtypes exist in close molecular proximity *in vivo* in whole animal models, or alternatively if these phenotypes result of signaling mechanisms that are independent of GPCR heteromeric formation. Detailed measurement of such molecular proximity *in vivo*, as well as the processes that control GPCR heteromerization in whole animal models, will require further study.

## LIMITATIONS, FUTURE DIRECTIONS, AND CONCLUDING REMARKS

Although a wealth of data from *in vitro* and in cellulo models have established the important role of GPCR heteromers in mediating precise and distinct roles in signaling cascades, their influence in the establishment of complex behavioral phenotypes remains to be fully elucidated. For instance, certain physiological and behavioral outcomes could conceivably be altered in the presence of peptides that according to *in vitro* or in cellulo assays disrupt GPCR heteromeric assembly. Similarly, the use of viral-mediated over-expression or transgenic animals could translate into animal models previous findings with receptor mutants that do not form heterocomplexes *in vitro* or in cellulo. However, a more precise understanding of such structural assembly obtained in rodent models will be necessary to fully define whether GPCR heteromers exist and function *in vivo*. Some of these strategies include the use of FRET (McGinty et al., 2011) or BRET (Dragulescu-Andrasi et al., 2011) imaging of protein-protein interactions in living mice.

An important challenge in the fields of GPCR research and molecular pharmacology is to develop an integrated understanding of how various mechanisms communicate with each other to ultimately orchestrate the formation of heteromeric complexes between some but not all GPCR subtypes. Potential mechanisms that are critical for this interaction specificity include specific pairs of residues that govern heteromeric formation, clustering of GPCRs in membrane microdomains, and crosstalk between receptors and a plethora of multidomain scaffolding proteins. Another important question to be addressed by future research is the molecular basis through which GPCR heteromers affect G protein function. For example, it has been shown that drugs that activate the G<sub>q/11</sub>-coupled 5-HT<sub>2A</sub> receptor induce both G<sub>q/11</sub>- and G<sub>i/o</sub>-dependent signaling in HEK293 cells co-expressing 5-HT<sub>2A</sub> and the G<sub>i/o</sub>-coupled mGlu2 receptor as a GPCR heteromer (Gonzalez-Maeso et al., 2008). Although findings in knockout mice suggest that co-expression of 5-HT<sub>2A</sub> and mGlu2 receptors is necessary to activate G<sub>q/11</sub> and G<sub>i/o</sub> by 5-HT<sub>2A</sub>

agonists in mouse frontal cortex membrane preparations (Fribourg et al., 2011), whether heteromeric formation is needed in living mice for this signaling crosstalk remains unknown. Similarly, more work is required both in cellulo and in animal models to solve whether  $G_{q/11}$  and  $G_{i/o}$  simultaneously or sequentially couple to the 5-HT<sub>2A</sub>-mGlu2 heteromeric receptor complex upon agonist binding to one of the two promoters.

Another significant limitation to our current understanding of GPCR heteromeric function is the lack of knowledge about physical stability of family A GPCR heteromers in animal models. Previous findings in HEK293 cells convincingly demonstrate that the  $\alpha 1B$ -adrenergic receptor forms higher-order oligomers, and that receptor oligomerization is required for receptor maturation and plasma membrane delivery (Lopez-Gimenez et al., 2007). On the other hand, results based on an experimental approach that recruits beta2-adrenergic receptors into artificial domains on the surface of living HEK293 cells suggest that the components of family A GPCR homomers interact transiently (Fonseca and Lambert, 2009; Gavalas et al., 2013). A similar conclusion has been reached using total internal reflection fluorescence microscopy (TIRFM) to visualize individual molecules in isolated CHO cells—the authors observed a transient association and dissociation of muscarinic M<sub>1</sub> receptor dimers in real time (Hern et al., 2010). Much further work is needed to characterize where along the pathway from synthesis to maturation and degradation do GPCR heteromers form. It also remains uncertain the stability of family A GPCR heteromers both *in vitro* and in whole animal models. Many studies examining homomeric GPCR interfaces report that residues of both TM1 and TM4 form symmetrical interfaces that lead to higher order species in heterologous expression systems (Guo et al., 2005, 2008), and this has been supported further by a number of recent crystal structures (Wu et al., 2012; Huang et al., 2013). However, it remains to be fully elucidated whether different homomeric and heteromeric organizations (e.g., squares and/or parallelograms) might exist in native tissue. These are all key questions that require further technical advances.

In conclusion, although a range of approaches has been applied and this has led to a general appreciation that GPCR heteromers affect receptor trafficking, pharmacology and function in cellulo, much more work is needed to probe the role of GPCR heteromerization *in vivo*. These advances in GPCR heteromeric research are now occurring at a rapid pace and promise to greatly contribute to the future of molecular medicine.

## ACKNOWLEDGMENT

Supported by R01MH084894 to Javier González-Maeso.

## REFERENCES

- Angers, S., Salahpour, A., Joly, E., Hilaliret, S., Chelsky, D., Dennis, M., et al. (2000). Detection of beta 2-adrenergic receptor dimerization in living cells using bioluminescence resonance energy transfer (BRET). *Proc. Natl. Acad. Sci. U.S.A.* 97, 3684–3689. doi: 10.1073/pnas.97.7.3684
- Baba, K., Benleulmi-Chaachoua, A., Journe, A. S., Kamal, M., Guillaume, J. L., Dussaud, S., et al. (2013). Heteromeric MT1/MT2 melatonin receptors modulate photoreceptor function. *Sci. Signal* 6, ra89. doi: 10.1126/scisignal.2004302
- Bourne, H. R., Sanders, D. A., and McCormick, F. (1990). The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348, 125–132. doi: 10.1038/348125a0
- Bouvier, M., and Hebert, T. E. (2014). CrossTalk proposal: weighing the evidence for Class A GPCR dimers, the evidence favours dimers. *J. Physiol.* 592, 2439–2441. doi: 10.1113/jphysiol.2014.272252
- Dragulescu-Andrasi, A., Chan, C. T., De, A., Massoud, T. F., and Gambhir, S. S. (2011). Bioluminescence resonance energy transfer (BRET) imaging of protein-protein interactions within deep tissues of living subjects. *Proc. Natl. Acad. Sci. U.S.A.* 108, 12060–12065. doi: 10.1073/pnas.1100923108
- Ferre, S., Casado, V., Devi, L. A., Filizola, M., Jockers, R., Lohse, M. J., et al. (2014). G protein-coupled receptor oligomerization revisited: functional and pharmacological perspectives. *Pharmacol. Rev.* 66, 413–434. doi: 10.1124/pr.113.008052
- Ferre, S., von Euler, G., Johansson, B., Fredholm, B. B., and Fuxe, K. (1991). Stimulation of high-affinity adenosine A2 receptors decreases the affinity of dopamine D2 receptors in rat striatal membranes. *Proc. Natl. Acad. Sci. U.S.A.* 88, 7238–7241. doi: 10.1073/pnas.88.16.7238
- Fonseca, J. M., and Lambert, N. A. (2009). Instability of a class A G protein-coupled receptor oligomer interface. *Mol. Pharmacol.* 75, 1296–1299. doi: 10.1124/mol.108.053876
- Fribourg, M., Moreno, J. L., Holloway, T., Provati, D., Baki, L., Mahajan, R., et al. (2011). Decoding the Signaling of a GPCR heteromeric complex reveals a unifying mechanism of action of antipsychotic drugs. *Cell* 147, 1011–1023. doi: 10.1016/j.cell.2011.09.055
- Gavalas, A., Lan, T. H., Liu, Q., Correa, I. R. Jr., Javitch, J. A., and Lambert, N. A. (2013). Segregation of family A G protein-coupled receptor protomers in the plasma membrane. *Mol. Pharmacol.* 84, 346–352. doi: 10.1124/mol.113.086868
- González-Maeso, J. (2011). GPCR oligomers in pharmacology and signaling. *Mol. Brain* 4:20. doi: 10.1186/1756-6606-4-20
- Gonzalez-Maeso, J., Ang, R. L., Yuen, T., Chan, P., Weisstaub, N. V., Lopez-Gimenez, J. F., et al. (2008). Identification of a serotonin/glutamate receptor complex implicated in psychosis. *Nature* 452, 93–97. doi: 10.1038/nature06612
- Gonzalez-Maeso, J., Weisstaub, N. V., Zhou, M., Chan, P., Ivic, L., Ang, R., et al. (2007). Hallucinogens recruit specific cortical 5-HT(2A) receptor-mediated signaling pathways to affect behavior. *Neuron* 53, 439–452. doi: 10.1016/j.neuron.2007.01.008
- Guo, W., Shi, L., Filizola, M., Weinstein, H., and Javitch, J. A. (2005). Crosstalk in G protein-coupled receptors: changes at the transmembrane homodimer interface determine activation. *Proc. Natl. Acad. Sci. U.S.A.* 102, 17495–17500. doi: 10.1073/pnas.0508950102
- Guo, W., Urizar, E., Kralikova, M., Mobarec, J. C., Shi, L., Filizola, M., et al. (2008). Dopamine D2 receptors form higher order oligomers at physiological expression levels. *EMBO J.* 27, 2293–2304. doi: 10.1038/emboj.2008.153
- Hanks, J. B., and Gonzalez-Maeso, J. (2013). Animal models of serotonergic psychedelics. *ACS Chem. Neurosci.* 4, 33–42. doi: 10.1021/cn300138m
- Hasbi, A., Perreault, M. L., Shen, M. Y., Zhang, L., To, R., Fan, T., et al. (2014). A peptide targeting an interaction interface disrupts the dopamine D1-D2 receptor heteromer to block signaling and function *in vitro* and *in vivo*: effective selective antagonism. *FASEB J.* doi: 10.1096/fj.14-254037. [Epub ahead of print].
- Hern, J. A., Baig, A. H., Mashanov, G. I., Birdsall, B., Corrie, J. E., Lazareno, S., et al. (2010). Formation and dissociation of M1 muscarinic receptor dimers seen by total internal reflection fluorescence imaging of single molecules. *Proc. Natl. Acad. Sci. U.S.A.* 107, 2693–2698. doi: 10.1073/pnas.0907915107
- Huang, J., Chen, S., Zhang, J. J., and Huang, X. Y. (2013). Crystal structure of oligomeric beta1-adrenergic G protein-coupled receptors in ligand-free basal state. *Nat. Struct. Mol. Biol.* 20, 419–425. doi: 10.1038/nsmb.2504
- Jones, K. A., Borowsky, B., Tamm, J. A., Craig, D. A., Durkin, M. M., Dai, M., et al. (1998). GABA(B) receptors function as a heteromeric assembly of the subunits GABA(B)R1 and GABA(B)R2. *Nature* 396, 674–679. doi: 10.1038/25348
- Kaupmann, K., Malitschek, B., Schuler, V., Heid, J., Froestl, W., Beck, P., et al. (1998). GABA(B)-receptor subtypes assemble into functional heteromeric complexes. *Nature* 396, 683–687. doi: 10.1038/25360
- Kern, A., Albarran-Zeckler, R., Walsh, H. E., and Smith, R. G. (2012). Apo-ghrelin receptor forms heteromers with DRD2 in hypothalamic neurons and is essential for anorexigenic effects of DRD2 agonism. *Neuron* 73, 317–332. doi: 10.1016/j.neuron.2011.10.038
- Lambert, N. A., and Javitch, J. A. (2014). CrossTalk opposing view: weighing the evidence for class A GPCR dimers, the jury is still out. *J. Physiol.* 592, 2443–2445. doi: 10.1113/jphysiol.2014.272997

- Lee, S. P., So, C. H., Rashid, A. J., Varghese, G., Cheng, R., Lanca, A. J., et al. (2004). Dopamine D1 and D2 receptor Co-activation generates a novel phospholipase C-mediated calcium signal. *J. Biol. Chem.* 279, 35671–35678. doi: 10.1074/jbc.M401923200
- Limbird, L. E., Meyts, P. D., and Lefkowitz, R. J. (1975). Beta-adrenergic receptors: evidence for negative cooperativity. *Biochem. Biophys. Res. Commun.* 64, 1160–1168. doi: 10.1016/0006-291X(75)90815-3
- Liu, X.-Y., Liu, Z.-C., Sun, Y.-G., Ross, M., Kim, S., Tsai, F.-F., et al. (2011). Unidirectional cross-activation of GRPR by MOR1D uncouples itch and analgesia induced by opioids. *Cell* 147, 447–458. doi: 10.1016/j.cell.2011.08.043
- Lopez-Gimenez, J. F., Canals, M., Padiani, J. D., and Milligan, G. (2007). The alpha1b-adrenoceptor exists as a higher-order oligomer: effective oligomerization is required for receptor maturation, surface delivery, and function. *Mol. Pharmacol.* 71, 1015–1029. doi: 10.1124/mol.106.033035
- Magalhaes, A. C., Holmes, K. D., Dale, L. B., Comps-Agrar, L., Lee, D., Yadav, P. N., et al. (2010). CRF receptor 1 regulates anxiety behavior via sensitization of 5-HT2 receptor signaling. *Nat. Neurosci.* 13, 622–629. doi: 10.1038/nn.2529
- McGinty, J., Stuckey, D. W., Soloviev, V. Y., Laine, R., Wylezinska-Arridge, M., Wells, D. J., et al. (2011). *In vivo* fluorescence lifetime tomography of a FRET probe expressed in mouse. *Biomed. Opt. Express* 2, 1907–1917. doi: 10.1364/BOE.2.001907
- Milligan, G. (2013). The prevalence, maintenance and relevance of GPCR oligomerization. *Mol. Pharmacol.* 84, 158–169. doi: 10.1124/mol.113.084780
- Milligan, G., and Bouvier, M. (2005). Methods to monitor the quaternary structure of G protein-coupled receptors. *FEBS J.* 272, 2914–2925. doi: 10.1111/j.1742-4658.2005.04731.x
- Moreno, J. L., Holloway, T., Albizu, L., Sealfon, S. C., and Gonzalez-Maeso, J. (2011). Metabotropic glutamate mGlu2 receptor is necessary for the pharmacological and behavioral effects induced by hallucinogenic 5-HT2A receptor agonists. *Neurosci. Lett.* 493, 76–79. doi: 10.1016/j.neulet.2011.01.046
- Moreno, J. L., Muguruza, C., Umali, A., Mortillo, S., Holloway, T., Pilar-Cuellar, E., et al. (2012). Identification of three residues essential for 5-HT2A-mGlu2 receptor heteromerization and its psychoactive behavioral function. *J. Biol. Chem.* 287, 44301–44319. doi: 10.1074/jbc.M112.413161
- Pei, L., Li, S., Wang, M., Diwan, M., Anisman, H., Fletcher, P. J., et al. (2010). Uncoupling the dopamine D1-D2 receptor complex exerts antidepressant-like effects. *Nat. Med.* 16, 1393–1395. doi: 10.1038/nm.2263
- White, J. H., Wise, A., Main, M. J., Green, A., Fraser, N. J., Disney, G. H., et al. (1998). Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature* 396, 679–682. doi: 10.1038/25354
- Whorton, M. R., Bokoch, M. P., Rasmussen, S. G., Huang, B., Zare, R. N., Kobilka, B., et al. (2007). A monomeric G protein-coupled receptor isolated in a high-density lipoprotein particle efficiently activates its G protein. *Proc. Natl. Acad. Sci. U.S.A.* 104, 7682–7687. doi: 10.1073/pnas.0611448104
- Wu, H., Wacker, D., Mileni, M., Katritch, V., Han, G. W., Vardy, E., et al. (2012). Structure of the human kappa-opioid receptor in complex with JDTic. *Nature* 485, doi: 10.1038/nature10939

**Conflict of Interest Statement:** The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 27 August 2014; paper pending published: 17 September 2014; accepted: 21 September 2014; published online: 09 October 2014.

Citation: González-Maeso J (2014) Family a GPCR heteromers in animal models. *Front. Pharmacol.* 5:226. doi: 10.3389/fphar.2014.00226

This article was submitted to *Neuropharmacology*, a section of the journal *Frontiers in Pharmacology*.

Copyright © 2014 González-Maeso. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.